

## ***Gingival Index***

### ***Procedure & Method Information***

*Name of Procedure/Method*    Gingival Index

*Abbreviation*    GI

*Purpose*    To assess the prevalence and severity of gingivitis.

*Year of Establishment*    1963

*Type of Procedure/Method*

*Developer(s)*    H. Loe and J. Silness

*Oral Condition Category*

### ***Background Information***

*Background Information*    The Gingival Index (GI) was developed by H. Loe and J. Silness in 1963 to assess the severity and prevalence of gingivitis by examining only the qualitative changes (i.e., severity of the lesion) of the gingival soft tissue. The GI does not take into account periodontal pocket depth, degrees of bone loss, or any other quantitative change of the periodontium (Loe, 1967).

The GI is one of the most widely accepted and used gingival bleeding indices due to its documented validity, reliability, and ease of use. However, even though the GI has demonstrated sufficient sensitivity to distinguish between groups with mild and severe gingivitis, it may not discriminate as well between the middle ranges (Burt and Eklund, 1999).

It can be used on all surfaces for the entire mouth or selected teeth (Burt and Eklund, 1999) or selected areas of all or selected teeth (Loe, 1967). It also can be used on large population groups or an individual.

*Changes Over Time*    None

### ***Procedure Method***

*Procedure Method*    To obtain the GI, the examiner first will need sufficient lighting, a mouth mirror, and probe. The teeth and gingiva (gums) also should be dried lightly with a blast of air and/or cotton rolls.

The buccal, lingual, mesial, and distal surface areas of six teeth are examined and scored according to the following criteria and scoring system (Loe and Silness, 1963). The six teeth that are evaluated are the upper right first molar, the upper right lateral incisor, the upper left first bicuspid, the lower left first molar, the lower left lateral incisor, and the lower right first bicuspid.

To calculate the GI for an individual, each of the four gingival areas (i.e., buccal, lingual, mesial, and distal) of the tooth is given a score from 0 to 3 as described in the following criteria (Loe and Silness, 1963). Then, the four scores from the gingival area are added and divided by 4 to give the GI for the tooth. Afterwards, the GI for the teeth are added and divided by the number of teeth examined (i.e., 6). In addition, the scores for the individual teeth (i.e., incisors, premolars, and molars) may be grouped to designate the GI for groups of teeth.

#### Criteria and Scoring for the Gingival Index (GI) (Loe and Silness, 1963)

0 = Absence of inflammation.

1 = Mild inflammation - slight change in color and little change in texture.

2 = Moderate inflammation - moderate glazing, redness, oedema, and hypertrophy. Bleeding on pressure.

3 = Severe inflammation - marked redness and hypertrophy. Tendency to spontaneous bleeding. Ulceration.

Source: Loe H, Silness J. Periodontal disease in pregnancy. I. Prevalence and severity. *Acta Odontol Scand.* 1963; 21:533-51.

Mild inflammation usually occurs from 0.1 to 1.0, where moderate inflammation occurs from 1.1 to 2.0, and severe inflammation scores between 2.1 and 3.0 (Loe and Silness, 1963; Loe, 1967).

#### *Established Modifications*

In 1967, Loe detailed the sequence of the examination procedure and slightly modified the procedure to include the entire dentition instead of six teeth (Marks et al., 1993). This detailed exam is as follows:

Using gentle probing pressure, the examination of all erupted teeth typically starts with the upper right second molar and continues over the midline to the upper left second molar. For teeth on the right side of the midline, the exam sequence is distal, buccal/labial, and mesial. On the left side, the exam sequence is mesial, buccal/labial, and distal. When the three surfaces (i.e., distal, buccal/labial, mesial) of all teeth have been scored, the lingual surfaces of all the upper or maxillary teeth are examined beginning with the upper left second molar.

For the lower or mandibular arch, the exam begins with the lower left second molar through to the right second molar. On the left side of the midline, the exam sequence is distal, buccal/labial, and mesial, and on the right side it is mesial, buccal/labial, and distal. Afterwards, all lingual surfaces are scored beginning with the left second molar.

However, according to research, analyses show no difference in the results when only one of the interproximal surfaces for a tooth is examined instead of both, although the score for the one interproximal surface was doubled to allow for the second interproximal surface and the total score for the tooth was divided by 4 (Loe, 1967).

Third molars or wisdom teeth are not examined or scored in the upper or lower arch. According to the developers, the scoring for the GI takes approximately 2 to 5 minutes under optimal conditions and with chair-side assistance.

The calculation for the GI of an individual remained the same except the maximum number of teeth examined in the denominator is 28 instead of 6.

#### Criteria and Scoring for the Gingival Index (GI) (Loe, 1967)

0 = Normal gingiva.

1 = Mild inflammation - slight change in color, slight edema. No bleeding on probing.

2 = Moderate inflammation - redness, edema, and glazing. Bleeding on probing.

3 = Severe inflammation - marked redness and edema. Ulceration. Tendency to spontaneous bleeding.

Source: Loe H. The Gingival Index, the Plaque Index, and the Retention Index Systems. J Periodontol, part II, 1967; 38(Suppl):610-6.

In 1983, I.B. Lamster, M.C. Alfano, M.C. Seiger, and J.M. Gordon introduced further modifications to the Gingival Index by changing the ordinal scoring system and the invasive examination procedure to a noninvasive approach. This modified version of the Gingival Index was recommended by Lobene in 1986 due to its increased sensitivity in the low region of the scoring scale and the elimination of the "bleeding on pressure" component (Lobene, 1986; Lobene, Mankodi, Ciano, Lamm, Charles, and Ross, 1989). For more information, see the Modified Gingival Index.

*Federal Survey  
Modifications*

None

### **References**

#### *References*

Textbooks, Manuals, and the Internet:

Burt BA, Eklund SA. Dentistry, Dental Practice, and the Community, 5th edition. Philadelphia: W.B. Saunders Company, 1999.

Northern Arizona University (1998). The Epidemiology of Periodontal Diseases. Retrieved September 13, 1999, from the World Wide Web:  
<http://www.nauonline.nau.edu/welcome/tdrive/dh418/lesson>.

Schlager S, Yuodelis R, Page RC, Johnson RH. Periodontal Diseases - Basis Phenomena, Clinical Management, and Occlusal and Restorative Interrelationships, 2nd ed. Philadelphia: Lea & Febiger, 1990.

Journals:

Ciano SG. Current status of indices of gingivitis. J Clin Periodontol. 1986 May;13(5):375-8.

Gordon JM, Lamster IB, Seiger MC. Efficacy of Listerine antiseptic in inhibiting the

development of plaque and gingivitis. J Clin Periodontol. 1985 Apr;12:697-704.

Lamster IB, Alfano MC, Seiger MC, Gordon JM. The effect of Listerine antiseptic on reduction of existing plaque and gingivitis. Clin Prev Dent. 1983;5:12-16.

Lobene RR, Mankodi SM, Ciancio SG, Lamm RA, Charles CH, Ross NM. Correlations among gingival indices: a methodology study. J Periodontol. 1989 Mar;60(3):159-62.

Lobene RR. Discussion: Current status of indices for measuring gingivitis. J Clin Periodontol. 1986 May;13(5):381-2.

Lobene RR, Weatherford T, Ross NM, Lamm RA, Menaker L. A modified gingival index for use in clinical trials. Clin Prev Dent. 1986 Jan-Feb;8(1):3-6.

Loe H. The Gingival Index, the Plaque Index, and the Retention Index Systems. J Periodontol, part II, 1967;38(Suppl):610-6.

Loe H, Silness J. Periodontal disease in pregnancy. I. Prevalence and severity. Acta Odont Scand. 1963; 21:533-551.

Marks RG, Magnusson I, Taylor M, Clouser B, Maruniak J, Clark WB. Evaluation of reliability and reproducibility of dental indices. J Clin Periodontol. 1993 Jan;20(1):54-8.

#### *Validity*

Nowicki D, Vogel RI, Melcer S, Deasy MJ. The gingival bleeding time index. J Periodontol. 1981 May;52(5):260-2.

#### *Reliability*

Kingman A, Loe H, Anerud A, Boysen H. Errors in measuring parameters associated with periodontal health and disease. J Periodontol. 1991 Aug;62(8):477-86.

Loe H. The Gingival Index, the Plaque Index, and the Retention Index Systems. J Periodontol, part II, 1967;38(Suppl):610-6.

Spolsky VW, Gornbein JA. Comparing measures of reliability for indices of gingivitis and plaque. J Periodontol. 1996 Sep;67(9):853-9.

---

### ***Listing of Publications with Surveys &***

#### *Surveys & Studies*

##### **International Surveys & Studies:**

Aass AM, Gjermo P. Comparison of oral hygiene efficacy of one manual and two electric toothbrushes. Acta Odontol Scand. 2000 Aug;58(4):166-70.

Cahen PM, Turlot JC, Frank RM, Obry-Musset AM. National survey of caries prevalence in 6-15-year-old children in France. J Dent Res. 1989 Jan;68(1):64-8.

Chava VK. An evaluation of the efficacy of a curved bristle and conventional toothbrush. A comparative clinical study. J Periodontol. 2000 May;71(5):785-9.

Meyers IA, McQueen MJ, Harbrow D, Seymour GJ. The surface effect of dentifrices. Aust Dent J. 2000 Jun;45(2):118-24.

Walter MH, Wolf BH, Schmidt AE, Boening KW, Koch R. Plaque, gingival health and post-operative sensitivity in titanium inlays and onlays: a randomized controlled clinical trial. J Dent. 2001 May;29(3):181-6.

#### United States Surveys & Studies:

Al-Banyan RA, Echeverri EA, Narendran S, Keene HJ. Oral health survey of 5-12-year-old children of National Guard employees in Riyadh, Saudi Arabia. Int J Paediatr Dent. 2000 Mar;10(1):39-45.

DePaola PF. The Massachusetts health survey. J Mass Dent Soc. 1983 Spring;32(1):10-1, 23-5.

Kingman A, Loe H, Anerud A, Boysen H. Errors in measuring parameters associated with periodontal health and disease. J Periodontol. 1991 Aug;62(8):477-86.

McClanahan SF, Bartizek RD, Biesbrock AR. Identification and consequences of distinct Loe-Silness gingival index examiner styles for the clinical assessment of gingivitis. J Periodontol. 2001 Mar;72(3):383-92.

Spolsky VW, Gornbein JA. Comparing measures of reliability for indices of gingivitis and plaque. J Periodontol. 1996 Sep;67(9):853-9.